

	商品名	商品説明	効能・効果	用法・用量	成分・分量	使用上の注意	保管および取扱い上の注意																									
日本語	龍角散せき止めカプセル コデポン	龍角散のせき止めカプセルは、せきに効く白い顆粒と、たんを出しやすくする赤い顆粒がダブルで作用し、眠れないほどのしつこいせきにも効果を発揮する鎮咳去痰薬です。	せき、たん	・次の量を水又は温湯で服用してください。 <table border="1"> <thead> <tr> <th>年齢</th> <th>1回量</th> <th>1日服用回数</th> </tr> </thead> <tbody> <tr> <td>大人(15歳以上)</td> <td>2カプセル</td> <td rowspan="2">3回</td> </tr> <tr> <td>8歳以上15歳未満</td> <td>1カプセル</td> </tr> <tr> <td>8歳未満</td> <td>服用しないこと</td> <td>—</td> </tr> </tbody> </table> <用法・用量に関する注意> (1)用法・用量を厳守してください。 (2)小児に服用させる場合には、保護者の指導監督のもとに服用させてください。 (3)カプセルの取り出し方: 右図のようにカプセルの入っているPTPシートの凸部を指先で強く押し、裏面のアルミ箔を破り、取り出して服用してください。(誤ってそのまま飲み込んだりすると食道粘膜に突き刺さるなど思わぬ事故につながります)	年齢	1回量	1日服用回数	大人(15歳以上)	2カプセル	3回	8歳以上15歳未満	1カプセル	8歳未満	服用しないこと	—	6カプセル(大人の1日量)中 <table border="1"> <thead> <tr> <th>成分</th> <th>含量</th> </tr> </thead> <tbody> <tr> <td>ジヒドロコデインリン酸塩</td> <td>18mg</td> </tr> <tr> <td>dℓ-メチルエフェドリン塩酸塩</td> <td>40mg</td> </tr> <tr> <td>ノスカピン</td> <td>60mg</td> </tr> <tr> <td>グアイフェネシン</td> <td>300mg</td> </tr> <tr> <td>クロルフェニラミンマレイン酸塩</td> <td>7.5mg</td> </tr> <tr> <td>無水カフェイン</td> <td>150mg</td> </tr> </tbody> </table> 添加物：パレイショデンプン、乳糖水和物、赤色2号、赤色102号、青色1号、ラウリル硫酸Na、亜硫酸水素Na、ゼラチン、セルロース、カルメロースCa、ヒプロメロース	成分	含量	ジヒドロコデインリン酸塩	18mg	dℓ-メチルエフェドリン塩酸塩	40mg	ノスカピン	60mg	グアイフェネシン	300mg	クロルフェニラミンマレイン酸塩	7.5mg	無水カフェイン	150mg	1. 次の人は服用前に医師、薬剤師又は登録販売者に相談してください。 (1) 医師の治療を受けている人。 (2) 妊婦又は妊娠していると思われる人。 (3) 高齢者 (4) 薬などによりアレルギー症状を起こしたことがある人。 (5) 次の症状のある人。高熱、排尿困難 (6) 次の診断を受けた人。心臓病、高血圧、糖尿病、緑内障、甲状腺機能障害 2. 服用後、次の症状があらわれた場合は副作用の可能性があるので、直ちに服用を中止し、この説明文書を持って医師、薬剤師又は登録販売者に相談してください。 <関係部位> <症状> 皮膚 発疹・発赤、かゆみ 消化器 吐き気・嘔吐、食欲不振 精神神経系 めまい 泌尿器 排尿困難 まれに下記の重篤な症状が起こることがあります。その場合は直ちに医師の診療を受けてください。 <症状の名称> 再生不良性貧血 <症状> 青あざ、鼻血、歯ぐきの出血、発熱、皮膚や粘膜が青白くみえる、疲労感、動悸、息切れ、気分が悪くなりくらっとする、血尿などがあらわれる <症状の名称> 無顆粒球症 <症状> 突然の高熱、さむけ、のどの痛みなどがあらわれる。 3. 服用後、次の症状があらわれることがあるので、このような症状の継続又は増強が見られた場合には、服用を中止し、医師、薬剤師又は登録販売者に相談してください。 便秘、口の渇き、眠気 4. 5～6回服用しても症状がよくなる場合は服用を中止し、この説明文書を持って医師、薬剤師又は登録販売者に相談してください。	(1)直射日光の当たらない湿気の少ない涼しい所に保管してください。 (2)小児の手の届かない所に保管してください。 (3)他の容器に入れ替えないでください(誤用の原因になったり品質が変わることがあります)。 (4)使用期限を過ぎた製品は服用しないでください。
年齢	1回量	1日服用回数																														
大人(15歳以上)	2カプセル	3回																														
8歳以上15歳未満	1カプセル																															
8歳未満	服用しないこと	—																														
成分	含量																															
ジヒドロコデインリン酸塩	18mg																															
dℓ-メチルエフェドリン塩酸塩	40mg																															
ノスカピン	60mg																															
グアイフェネシン	300mg																															
クロルフェニラミンマレイン酸塩	7.5mg																															
無水カフェイン	150mg																															
	Name	Product Description	Efficacy	Dosing and Administration	Ingredients	Warning	Warning in preservation																									
英語	Ryukakusan Cough Medicine(Capsule) "Kodepon"	Ryukakusan Cough Medicine(Capsule) has white and red granules. The white ones are to suppress coughs and the red ones clear the phlegm. The synergy effects of two ingredients help of relief from consistent coughs.	Cough, phlegm	Take the following doses with water. <table border="1"> <thead> <tr> <th>Age</th> <th>Dosage</th> <th>Daily usage</th> </tr> </thead> <tbody> <tr> <td>Adult(over 15)</td> <td>2 capsules</td> <td rowspan="2">3 times</td> </tr> <tr> <td>8 ~ under 15</td> <td>1 capsule</td> </tr> <tr> <td>under 8</td> <td>Do not use</td> <td>—</td> </tr> </tbody> </table> <Dosing and Administration.> 1. Please follow the dosing and administration. 2. For small children, take under the adults' guidance and supervision. 3. When taking out the capsule, push the convey side of the PTP sheet where the capsule is inserted just as the diagram labelled on right. (Swallowing without pushing it out could lead to accidents such as damaging the esophageal mucous membrane)	Age	Dosage	Daily usage	Adult(over 15)	2 capsules	3 times	8 ~ under 15	1 capsule	under 8	Do not use	—	6 capsules(Adult daily usage) <table border="1"> <thead> <tr> <th>Active Ingredient</th> <th>Dosage</th> </tr> </thead> <tbody> <tr> <td>Dihydrocodeine Phosphate</td> <td>18mg</td> </tr> <tr> <td>dℓ-Methylephedrine Hydrochloride</td> <td>40mg</td> </tr> <tr> <td>Noscapine</td> <td>60mg</td> </tr> <tr> <td>Guaifenesin</td> <td>300mg</td> </tr> <tr> <td>Chlorpheniramine Maleate</td> <td>7.5mg</td> </tr> <tr> <td>Anhydrous Caffeine</td> <td>150mg</td> </tr> </tbody> </table> Additives: Potato Starch, Lactose, Food Red No.2, Food Red No.102, Food Blue No.1, Sodium Lauryl Sulfate, Sodium Bisulfite, Gelatin, Cellulose, Carmellose Calcium, Hypromellose	Active Ingredient	Dosage	Dihydrocodeine Phosphate	18mg	dℓ-Methylephedrine Hydrochloride	40mg	Noscapine	60mg	Guaifenesin	300mg	Chlorpheniramine Maleate	7.5mg	Anhydrous Caffeine	150mg	1. Please consult with your physician, pharmacist or registered sales clerks. (1)Currently under a doctor's care. (2)People who are pregnant or having a possibility of pregnancy. (3)Elderly people (4)Experienced any allergy reactions by taking medication. (5)Having a high fever, dysuria (6)People diagnosed as follows. Heart disease, Hypertension, Diabetes, Glaucoma, Dysfunction thyroid 2. If you have the following reactions to the product, discontinue to use immediately. Take this product instruction to consult with your physician, pharmacist, or registered sales clerks. <Symptoms> Skin: Rash, itchiness, skin reddening Digestive system: nausea, vomiting, inappetence anorexia Nervous system: Dizziness Urinary system: Dysuria In case of examined as a serious symptom labelled below, contact and receive treatment from a doctor immediately. <Name of the symptom> anaplastic anemia <Symptom> Bruise, nasal bleeding, gum bleeding, fever, pale skin, feebleness, heart palpitation, short breath, feeling dizzy, bloody urine. <Name of the symptom> agranulocytosis <Symptom> Sudden high fever, shivering, throat pain 3. After taking this medicine, the following symptoms may occur. If the symptoms continue or enhance stop taking it immediately. Consult with your physician, pharmacist, or registered sales clerks. constipation, oral dryness, sleepiness 4. If your condition persists after usage of 5 to 6 times, stop taking this medicine. Please take this product instruction to show, and consult with your physician, pharmacist, or registered sales clerks.	(1) Store them in a cool, low humidity area avoiding from a direct sunlight. (2) Keep it away from small children. (3) Don't replace the product to a different container. (It may cause accidental ingestion and quality deterioration.) (4) Do not take expired products.
Age	Dosage	Daily usage																														
Adult(over 15)	2 capsules	3 times																														
8 ~ under 15	1 capsule																															
under 8	Do not use	—																														
Active Ingredient	Dosage																															
Dihydrocodeine Phosphate	18mg																															
dℓ-Methylephedrine Hydrochloride	40mg																															
Noscapine	60mg																															
Guaifenesin	300mg																															
Chlorpheniramine Maleate	7.5mg																															
Anhydrous Caffeine	150mg																															