

	商品名	商品説明	効能・効果	用法・用量	成分・分量	使用上の注意	保管および取扱い上の注意																														
日本語	龍角散せき止め錠	こどもでも大人でも、せきが止まらない時は本当につらいものです。つらいせきがずっと、睡眠もさまたげられ体力も消耗しますので、できる限り早く止めることが大切です。龍角散せき止め錠は、たんの排出を容易にするブロムヘキシン塩酸塩(去痰剤)、アレルギー性のせきに効果のあるクレマスチンフマル酸塩(抗ヒスタミン剤)、せき中枢を抑制してせきを鎮めるジヒドロコデインリン酸塩(鎮咳剤)などを配合して、すみやかにせきやたんを改善するお薬です。さらにクレマスチンフマル酸塩には、他の抗ヒスタミン剤と比較してねむけや口のかわきなどの副作用が少ないという特徴があります。	せき、たん	・次の量を食後に服用してください。 <table border="1"> <thead> <tr> <th>年齢</th> <th>1回量</th> <th>1日服用回数</th> </tr> </thead> <tbody> <tr> <td>大人(15歳以上)</td> <td>3錠</td> <td rowspan="3">3回</td> </tr> <tr> <td>11歳以上15歳未満</td> <td>2錠</td> </tr> <tr> <td>5歳以上11歳未満</td> <td>1錠</td> </tr> </tbody> </table> 5歳未満 服用しないこと <用法・用量に関する注意> (1)用法・用量を厳守してください。 (2)小児に服用させる場合には、保護者の指導監督のもとに服用させてください。	年齢	1回量	1日服用回数	大人(15歳以上)	3錠	3回	11歳以上15歳未満	2錠	5歳以上11歳未満	1錠	9錠(大人の1日量)中 <table border="1"> <thead> <tr> <th>成分</th> <th>含量</th> </tr> </thead> <tbody> <tr> <td>ブロムヘキシン塩酸塩</td> <td>12mg</td> </tr> <tr> <td>クレマスチンフマル酸塩</td> <td>1.34mg</td> </tr> <tr> <td>ジヒドロコデインリン酸塩</td> <td>30mg</td> </tr> <tr> <td>dℓ-メチルエフェドリン塩酸塩</td> <td>75mg</td> </tr> <tr> <td>ノスカピン</td> <td>60mg</td> </tr> <tr> <td>無水カフェイン</td> <td>90mg</td> </tr> <tr> <td>カンゾウ乾燥エキス</td> <td>105mg(819mg)</td> </tr> <tr> <td>キキョウ乾燥エキス末</td> <td>120mg(600mg)</td> </tr> <tr> <td>セネガ乾燥エキス</td> <td>36mg(600mg)</td> </tr> </tbody> </table> ( ) : 原生薬換算量 添加物: ヒドロキシプロピルセルロース、CMC-Ca、ケイ酸Aℓ、乳糖水和物、セルロース、メタケイ酸アルミン酸Mg、ステアリン酸Mg、ヒプロメロース、マクロゴール、酸化チタン、カルナウバロウ、黄色5号	成分	含量	ブロムヘキシン塩酸塩	12mg	クレマスチンフマル酸塩	1.34mg	ジヒドロコデインリン酸塩	30mg	dℓ-メチルエフェドリン塩酸塩	75mg	ノスカピン	60mg	無水カフェイン	90mg	カンゾウ乾燥エキス	105mg(819mg)	キキョウ乾燥エキス末	120mg(600mg)	セネガ乾燥エキス	36mg(600mg)	[してはいけないこと] 1. 次の人は服用前に医師、薬剤師又は登録販売者に相談してください。 (1) 医師の治療を受けている人。 (2) 妊婦又は妊娠していると思われる人。 (3) 高齢者 (4) 薬などによりアレルギー症状を起こしたことがある人。 (5) 次の症状のある人。高熱、排尿困難 (6) 次の診断を受けた人。心臓病、高血圧、糖尿病、緑内障、甲状腺機能障害。 2. 服用後、次の症状があらわれた場合は副作用の可能性があるので、直ちに服用を中止し、この説明文書を持って医師、薬剤師又は登録販売者に相談してください。 <関係部位> <症状> 皮膚 発疹・発赤、かゆみ 消化器 吐き気・嘔吐、食欲不振 精神神経系 めまい・興奮・けいれん 泌尿器 排尿困難 まれに下記の重篤な症状が起こることがあります。その場合は直ちに医師の診療を受けてください。 <症状の名称> ショック (アナフィラキシー) <症状> 服用後すぐに、皮膚のかゆみ、じんましん、声のかすれ、くしゃみ、のどのかゆみ、息苦しさ、動悸、意識の混濁などがあらわれる。 3. 服用後、次の症状があらわれることがあるので、このような症状の持続又は増強が見られた場合には、服用を中止し、医師、薬剤師又は登録販売者に相談してください。 便秘、口のかわき、眠気 4. 5～6回服用しても症状がよくならない場合は服用を中止し、この説明文書を持って医師、薬剤師又は登録販売者に相談してください。	(1) 直射日光の当たらない湿気の少ない涼しい所に密栓して保管してください。 (2) 小児の手の届かない所に保管してください。 (3) 他の容器に入れ替えないでください(誤用の原因になったり品質が変わることがあります)。 (4) 使用期限を過ぎた製品は服用しないでください。 (5) 水分が錠剤に付着しますと表面のフィルム層の一部が溶けて斑点を生じることがありますので、誤って水滴を落としたり、ぬれた手で触れないようにしてください。  2011年10月改訂 (記載要領変更に伴う改正)
年齢	1回量	1日服用回数																																			
大人(15歳以上)	3錠	3回																																			
11歳以上15歳未満	2錠																																				
5歳以上11歳未満	1錠																																				
成分	含量																																				
ブロムヘキシン塩酸塩	12mg																																				
クレマスチンフマル酸塩	1.34mg																																				
ジヒドロコデインリン酸塩	30mg																																				
dℓ-メチルエフェドリン塩酸塩	75mg																																				
ノスカピン	60mg																																				
無水カフェイン	90mg																																				
カンゾウ乾燥エキス	105mg(819mg)																																				
キキョウ乾燥エキス末	120mg(600mg)																																				
セネガ乾燥エキス	36mg(600mg)																																				
英語	Ryukakusan Cough Medicine(Tablet)	From children to adults, it is harsh getting diagnosed with a tough cough. Continuation of a tough cough can lead to lack of sleep making the body exhausted, requiring an immediate cure. Ryukakusan Cough Medicine(Tablet) contains bromhexine hydrochloride(expectorant) enhancing phlegm extrusion, clemastine fumarate(antihistaminic) healing allergic coughs, dihydrocodeine phosphate stopping coughs from calming down the cough centrum, it is a medicine to improve throat and phlegm quickly. Comparing to other antihistaminic medicines, clemastine fumarate contains low side effects of sleepiness and mouth dryness.	Cough, phlegm	Take the following doses after meals. <table border="1"> <thead> <tr> <th>Age</th> <th>Usage</th> <th>Daily usage</th> </tr> </thead> <tbody> <tr> <td>Adult(over 15)</td> <td>3 tablets</td> <td rowspan="3">3 times</td> </tr> <tr> <td>11 ~ under 15</td> <td>2 tablets</td> </tr> <tr> <td>5 ~ under 11</td> <td>1 tablet</td> </tr> </tbody> </table> under 5 Do not use <Dosing and Administration.> 1. Please follow the dosing and administration. 2. For small children, take under the adults' guidance and supervision.	Age	Usage	Daily usage	Adult(over 15)	3 tablets	3 times	11 ~ under 15	2 tablets	5 ~ under 11	1 tablet	9 tablets(Adult daily usage) <table border="1"> <thead> <tr> <th>Active Ingredient</th> <th>Dosage</th> </tr> </thead> <tbody> <tr> <td>Bromhexine Hydrochloride</td> <td>12mg</td> </tr> <tr> <td>Clemastine Fumarate</td> <td>1.34mg</td> </tr> <tr> <td>Dihydrocodeine Phosphate</td> <td>30mg</td> </tr> <tr> <td>dℓ-Methylephedrine Hydrochloride</td> <td>75mg</td> </tr> <tr> <td>Noscapine</td> <td>60mg</td> </tr> <tr> <td>Anhydrous Caffeine</td> <td>90mg</td> </tr> <tr> <td>Glycyrrhiza dry Extract</td> <td>105mg(819mg)</td> </tr> <tr> <td>Powdered Platycodon dry Extract</td> <td>120mg(600mg)</td> </tr> <tr> <td>Senega dry Extract</td> <td>36mg(600mg)</td> </tr> </tbody> </table> ( ) : Original Herbal Reduced Quantity Additives: Hydroxypropylcellulose, Carmellose Calcium, Aluminium Silicate, Lactose, Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate, Hypromellose, Macrogol, Titanium Oxide, Carnauba Wax, Food Yellow No.5	Active Ingredient	Dosage	Bromhexine Hydrochloride	12mg	Clemastine Fumarate	1.34mg	Dihydrocodeine Phosphate	30mg	dℓ-Methylephedrine Hydrochloride	75mg	Noscapine	60mg	Anhydrous Caffeine	90mg	Glycyrrhiza dry Extract	105mg(819mg)	Powdered Platycodon dry Extract	120mg(600mg)	Senega dry Extract	36mg(600mg)	1. Please consult with your physician, pharmacist or registered sales clerks if you are: (1) Currently under a doctor's care. (2) People who are pregnant or having a possibility of pregnancy. (3) Elderly people (4) Experienced any allergy reactions by taking medication. (5) Having a high fever, dysuria (6) People diagnosed as follows. Heart disease, Hypertension, Diabetes, Glaucoma, Dysfunction thyroid 2. If you have the following reactions to the product, discontinue to use immediately. Take this product instruction to consult with your physician, pharmacist, or registered sales clerks. <Symptoms> Skin: Rash, skin reddening, itchiness Digestive system: nausea, vomiting, inappetence anorexia Nervous system: Dizziness Urinary system: Dysuria In case of examined as a serious symptom labelled below, contact and receive treatment from a doctor immediately. <Name of the symptom> anaphylactic shock <Symptom> Soon after use, having symptoms such as skin itchiness, hives, hoarse voices, sneezing, throat itchiness, dyspnea, heart palpitation, obnubilation, 3. After taking this medicine, the following symptoms may occur. If the symptoms continue or enhance stop taking it immediately. Consult with your physician, pharmacist, or registered sales clerks. constipation, oral dryness, sleepiness 4. If your condition persists after usage of 5 to 6 times, stop taking this medicine. Please take this product instruction to show, and consult with your physician, pharmacist, or registered sales clerks.	(1) Store them in a cool, low humidity area avoiding from a direct sunlight. (2) Keep it away from small children. (3) Don't replace the product to a different container. (It may cause accidental ingestion and quality deterioration.) (4) Do not take expired products. (5) Drop of water on the surface may cause the spotty appearance of the film coating. Do not touch with wet hands or mistakenly spill water on it.  Revised on October, 2011 (Description change)
Age	Usage	Daily usage																																			
Adult(over 15)	3 tablets	3 times																																			
11 ~ under 15	2 tablets																																				
5 ~ under 11	1 tablet																																				
Active Ingredient	Dosage																																				
Bromhexine Hydrochloride	12mg																																				
Clemastine Fumarate	1.34mg																																				
Dihydrocodeine Phosphate	30mg																																				
dℓ-Methylephedrine Hydrochloride	75mg																																				
Noscapine	60mg																																				
Anhydrous Caffeine	90mg																																				
Glycyrrhiza dry Extract	105mg(819mg)																																				
Powdered Platycodon dry Extract	120mg(600mg)																																				
Senega dry Extract	36mg(600mg)																																				