

商品名	商品説明	効能・効果	用法・用量	成分・分量	使用上の注意	保管および取扱い上の注意
龍角散鼻炎朝タカセル	龍角散鼻炎朝タカセルは、抗アレルギー作用の持続時間が長いクレマチンマル酸塩と生薬成分サイシンエキスを同時に配合しました。花粉症などのアレルギー性鼻炎によるくしゃみや鼻みず、鼻づまり、なみだ目、のどの痛み、頭痛(頭が重い)	急性鼻炎、アレルギー性鼻炎又は副鼻腔炎による次の諸症状の緩和： くしゃみ、鼻みず(鼻汁過多)、鼻づまり、なみだ目、のどの痛み、頭痛(頭が重い)	次の量を、朝・夕に水又は温湯で服用してください。  年齢   1回量   1日服用回数 大人(15歳以上)   1カプセル   2回  15歳未満   服用しないでください	成分 クレマチンマル酸塩 1.34mg 主なほたらき 抗ヒスタミン作用により、くしゃみ、鼻みず、鼻づまりなどを持続的におさえます。  グリチルリチン酸二カリウム 50mg 抗炎症作用、抗アレルギー作用により、くしゃみ、鼻みずなどをおさえます。  サイシンエキス(原生薬換算量) 60mg(600mg) 抗アレルギー作用により、くしゃみ、鼻みずなどをおさえます。  添加物：白糖、エチルセルロース、ヒプロメロース、マクロゴール、タルク、ゼラチン、三酸化鉄、酸化チタン、ラウリル硫酸Na、青色1号	【してはいけないこと】 ① 本剤を使用している間は、次のいずれの医薬品も使用しないでください。 他の鼻炎用内服薬、抗ヒスタミン剤を含有する内服薬等(かぜ薬、鎮咳去痰薬、乗酔薬、アレルギー用薬等) ② 服用後、乗物又は機械類の運転操作をしないでください。(眠気等があらわれることがあります。) ③ 長期運用しないでください。  【相談すること】 1. 次の人は服用前に医師、薬剤師又は登録販売者に相談してください。 (1) 医師の治療を受けている人。 (2) 妊婦又は妊娠していると思われる人。 (3) 授乳中の人 (3) 高齢者 (4) 薬などによりアレルギー症状を起こしたことがある人。 (5) 次の症状のある人。高熱、排尿困難、むくみ (6) 次の診断を受けた人。緑内障、心臓病、高血圧、腎臓病。  2. 服用後、次の症状があらわれた場合は副作用の可能性があるので、直ちに服用を中止し、この説明文書を持って医師、薬剤師又は登録販売者に相談してください。 <関係部位> <症状> 皮膚 発疹・発赤、かゆみ 消化器 吐き気・嘔吐、食欲不振 精神神経系 興奮、痙攣 泌尿器 排尿困難 まれに下記の重篤な症状が起こることがあります。その場合は直ちに医師の診療を受けてください。  <症状の名称> 肝機能障害  <症状> 発熱、かゆみ、発疹、黄疽(皮膚や白目が黄色くなる)、褐色尿、全身のだるさ、食欲不振等があらわれる。  <症状の名称> 偽アルドステロン症、ミオパチー  <症状> 手足のだるさ、しびれ、つっぱり感やこわばりに加えて、脱力感、筋肉痛があらわれ、徐々に強くなる。  3. 服用後、次の症状があらわれることがあるので、このような症状の持続又は増強が見られた場合には、服用を中止し、医師、薬剤師又は登録販売者に相談してください。 口のかき、眠気  4. 5～6日間服用しても症状がよくならない場合は服用を中止し、この説明文書を持って医師、薬剤師又は登録販売者に相談してください。	(1)直射日光の当たらない湿気の少ない涼しい所に保管してください。 (2)小児の手の届かない所に保管してください。 (3)他の容器に入れ替えないでください(誤用の原因になったり品質が変わることがあります)。 (4)使用期限を過ぎた製品は服用しないでください。  2012年7月改訂 (記載要領変更に伴う改正)

Name	Product Description	Efficacy	Dosing and Administration	Ingredients	Warning	Warning in preservation
Ryukakusan Sinus Day and Night Capsule	Ryukakusan's Sinus Day and Night Capsule contains clemastine fumarate and crude drugs asiasarum extract, which have a long lasting effect for the allergic reaction. It eases allergic nasal catarrh sneezes and snivels caused from pollenosis for a long time. Clemastine fumarate contains less side effects of sleepiness, or dry mouth. The crude drug contains anti allergic effects. It eases the sneezes and allergic symptoms gently. Take one capsule twice a day, day and night. It lasts for 24 hours.	Easing the following symptoms caused from acute rhinitis, allergic rhinitis, or sinusitis: Sneezing, snivel(mucus nasal increased), nasal congestion, epiphora, sore throat, headache(heaviness of head)	Take the following dosage day and night using water.  Age   Usage   Daily usage Adult(over 15)   1 capsule   2 times  under 15   Do not use	2 capsules(Adult daily usage)  Active Ingredient   Dosage   Main effect Clemastine Fumarate   1.34mg   Antihistaminic action suppresses snivel, nasal congestion continuously.  Dipotassium Glycyrrhizinate   50mg   Anti-inflammatory action and anti allergic action suppresses sneezing, snivel, etc.  Asiasarum Extract(Original Herbal Reduced Quantity)   60mg(600mg)   Anti allergic action suppresses sneezing, snivel, etc.  Additives: Sucrose, Ethylcellulose, Hypromellose, Macrogol, Talc, Gelatin, Red Ferric Oxide, Titanium Oxide, Sodium Lauryl Sulfate, Food Blue No.1	【Side effects or any adverse events may occur by not following the direction.】 ① Do not take following medicines while using this medicine. Taking other sinus oral medicine, oral medicines containing antihistaminic. (Cold medicine, cough medicine, motion sickness drug, allergy medicine) ② Do not drive a motor vehicle or operate machinery when using this product. It may cause drowsiness. ③ Do not use for extended period.  【What to ask】 1. Please consult with your physician, pharmacist or registered sales clerks if you are: (1)Currently under a doctor's care. (2)People who are pregnant or having a possibility of pregnancy. (3)Elderly people (4)Experienced any allergy reactions by taking medication. (5)Having a high fever, dysuria, oedema. (6)People diagnosed as follows: Glaucoma, Heart Disease, Hypertension, Kidney Disease.  2. If you have the following reactions to the product, discontinue to use immediately. Take this product instruction to consult with your physician, pharmacist, or registered sales clerks.  <Symptoms> Skin: Rash, skin reddening, itchiness Digestive system: nausea, vomiting, inappetence anorexia Nervous system: Dizziness Urinary system: Dysuria In case of having a serious symptom labelled below, contact and receive treatment from a doctor immediately.  <Name of the symptom> Liver function disorder <Symptom> Fever, Itchiness, Exanthema, Jaundice(Skin and white of eye becomes yellow), hemosiderinuria, sluggishness, inappetence anorexia, etc.  <Name of the symptom> Pseudoaldosteronism, myopathy <Symptom> Starting to have tiredness, numbness, stiffness, feeling weak, muscle pain, and gradually symptoms get severe.  3. After taking this product, the following symptoms may occur. If the symptoms continue or enhance stop taking it immediately. Consult with your physician, pharmacist, or registered sales clerks. oral dryness, sleepiness  4. If your condition persists more than 5 to 6 days, stop taking this medicine. Please take this product instruction to show, and consult with your physician, pharmacist, or registered sales clerks.	(1) Store them in a cool, low humidity area avoiding from a direct sunlight. (2) Keep it away from small children. (3) Don't replace the product to a different container. (It may cause accidental ingestion and quality deterioration.) (4) Don't take expired products.  Revised on July 2012 (Description change)

日本語

英語